


## Lecture Series 3 Macromolecules: Their Structure and Function


### Biological Substances found in Living Tissues



### A. Lipids: Water-Insoluble Molecules

- Lipids can form large biological molecules, but these aggregations are NOT chemically polymers because individual units are not linked by covalent bonds.
- Share the common trait of being hydrophobic.


### The role of hydrocarbons in fats





### A. Lipids: Water-Insoluble Molecules

- Fats and oils are composed of three fatty acids covalently bonded to a glycerol molecule by ester linkages.
- Fats and oils function to efficiently store energy.

### Synthesis of a Triglyceride









**A. Lipids: Water-Insoluble Molecules**

- Carotenoids trap light energy in green plants.  $\beta$ -Carotene can be split to form vitamin A, a lipid vitamin.



**A. Lipids: Water-Insoluble Molecules**

- Some lipids are steroids and function as hormones. Cholesterol is synthesized by the liver and has a role in some cell membranes, as well as in the digestion of other fats.
- Some lipids function as vitamins, required for normal functioning, must be acquired from the diet.




## B. Macromolecules: Giant Polymers

- Macromolecules have specific three-dimensional shapes. Different functional groups give local sites on macromolecules specific properties.
- Monomers are joined by condensation reactions. Hydrolysis reactions break polymers into monomers.


### 3.1 The Building Blocks of Organisms

| MONOMER        | SIMPLE POLYMER          | COMPLEX POLYMER |
|----------------|-------------------------|-----------------|
| Amino acid     | Peptide or oligopeptide | Polypeptide     |
| Nucleotide     | Oligonucleotide         | Nucleic acid    |
| Monosaccharide | Oligosaccharide         | Polysaccharide  |

The synthesis and breakdown of polymers




Condensation or Dehydration reactions




Hydrolysis reactions

The synthesis and structure of a fat, or triacylglycerol




Glycerol  
(a) Dehydration synthesis







## C. Carbohydrates: Sugars and Sugar Polymers

- All carbohydrates contain carbon bonded to H and OH groups.  $[\text{CH}_2\text{O}]_N$
- Hexoses are monosaccharides that contain six carbon atoms.
- Monosaccharides are simple sugars.
  - ◆ Can be used for fuel.
  - ◆ Can be converted into other organic molecules.
  - ◆ Can be combined into polymers.

Various forms of Glucose






May be linear, but can form rings.



### C. Carbohydrates: Sugars and Sugar Polymers



- Glycosidic linkages may have either  $\alpha$  or  $\beta$  orientation in space. They covalently link monosaccharides into larger units.





### C. Carbohydrates: Sugars and Sugar Polymers

- Cellulose, a polymer, is formed by glucose units linked by  $\beta$ -glycosidic linkages between carbons 1 and 4.



- Cellulose is difficult to digest
  - Cows have microbes in their stomachs to facilitate this process


## C. Carbohydrates: Sugars and Sugar Polymers

- Starches are formed by  $\alpha$ -glycosidic linkages between carbons 1 and 4 and are distinguished by amount of branching through glycosidic bond formation at carbon 6.
- Glycogen contains  $\alpha$ -1,4 glycosidic linkages and is highly branched.


## Glycosidic Linkages

### (a) Molecular structure


Starch and glycogen



## Storage polysaccharides



## Starch and cellulose molecular models




Benjamin Cummings

## C. Carbohydrates: Sugars and Sugar Polymers


- Chemically modified monosaccharides include the sugar phosphates and amino sugars. A derivative of the amino sugar glucosamine polymerizes to form the polysaccharide chitin.

### (a) Sugar phosphate




Modified Sugars

### (b) Amino sugars



### (c) Chitin



- Chitin, another important structural polysaccharide
  - ◆ Is found in the exoskeleton of arthropods
  - ◆ Can be used as surgical thread



Chitin forms the exoskeleton of arthropods. This cicada is molting, shedding its old exoskeleton and emerging in adult form.


Chitin is used to make a strong and flexible surgical thread that decomposes after the wound or incision heals.

## D. Proteins: Amazing Polymers of Amino Acids

- Functions of proteins include support, protection, catalysis, transport, defense, regulation, and movement. They sometimes require an attached prosthetic group.
- Twenty amino acids are found in proteins. Each consists of an amino group, a carboxyl group, a hydrogen, and a side chain bonded to the  $\alpha$  carbon atom.


### 3.2 Twenty Amino Acids Found in Proteins

#### A. Amino acids with electrically charged side chains




### 3.2 Twenty Amino Acids Found in Proteins

#### B. Amino acids with polar but uncharged side chains




#### C. Special cases



### 3.2 Twenty Amino Acids Found in Proteins


#### D. Amino acids with hydrophobic side chains



## D. Proteins: Amazing Polymers of Amino Acids



- Side chains of amino acids may be charged, polar, or hydrophobic. SH groups can form disulfide bridges.





## D. Proteins: Amazing Polymers of Amino Acids

- Amino acids are covalently bonded together by peptide linkages.





## D. Proteins: Amazing Polymers of Amino Acids

- Polypeptide chains of proteins are folded into specific three-dimensional shapes. Primary, secondary, tertiary, and quaternary structures are possible.

## D. Proteins: Amazing Polymers of Amino Acids



- The primary structure of a protein is the sequence of amino acids bonded by peptide linkages.
- Secondary structures are maintained by hydrogen bonds between atoms of the amino acid residues.






## D. Proteins: Amazing Polymers of Amino Acids


- The tertiary structure is generated by bending and folding of the polypeptide chain. This results from interactions between amino acids and R groups.
- The quaternary structure is the arrangement of polypeptides in a single functional unit consisting of more than one polypeptide subunit.



## D. Proteins: Amazing Polymers of Amino Acids


- Weak chemical interactions are important in the binding of proteins to other molecules.
- Proteins denatured by heat, acid, or chemicals lose tertiary and secondary structure and biological function.





### D. Proteins: Amazing Polymers of Amino Acids

- Chaperonins assist protein folding by preventing binding to inappropriate ligands.
- They also help to shape proteins with special needs regarding hydrophobic and hydrophilic interactions.




### E. Nucleic Acids: Informational Macromolecules

- In cells, DNA is the hereditary material. DNA and RNA play roles in protein formation.

### E. Nucleic Acids: Informational Macromolecules

- Nucleic acids are polymers of nucleotides consisting of a phosphate group, a sugar, and a nitrogen-containing base. The DNA bases are adenine, guanine, cytosine, and thymine. In RNA uracil substitutes for thymine and ribose substitutes for deoxyribose.



### 3.3 Distinguishing RNA from DNA



| NUCLEIC ACID | SUGAR       | BASES                                     |
|--------------|-------------|---|
| RNA          | Ribose      | Adenine<br>Cytosine<br>Guanine<br>Uracil  |
| DNA          | Deoxyribose | Adenine<br>Cytosine<br>Guanine<br>Thymine |

## E. Nucleic Acids: Informational Macromolecules


- In the nucleic acids, bases extend from a sugar–phosphate backbone using the phosphodiester linkage.
- DNA and RNA information resides in their base sequences.

## E. Nucleic Acids: Informational Macromolecules

- RNA is single-stranded.
- DNA is a double-stranded helix with complementary, hydrogen-bonded base pairing between adenine and thymine and guanine and cytosine. The two strands run in opposite 5' to 3' directions.



DNA structure: The double helix



## E. Nucleic Acids: Informational Macromolecules

- Comparing the DNA base sequences of different living species provides information on evolutionary relatedness.
- This is called molecular phylogeny.

## F. The Interactions of Macromolecules

- Glycoproteins contain an oligosaccharide "label" that directs the protein to the proper cell destination. The carbohydrate groups of glycolipids are on the cell's outer surface, serving as recognition signals.
- An example of emergent properties where greater complexity is exhibited.