
Lecture Series 6 DNA and Its Role in Heredity


Reading Assignments

- Read Chapter 5
 DNA and Chromosomes
- Read Chapter 6
 DNA Replication, Repair & Replication (skim pages 215 to 222 on Recombination)
- Read Chapter 10
 Pages 331 to 333 and 347 to 351
 (only sections on Sequencing and PCR)

A. DNA: The Genetic Material


- In addition to circumstantial evidence, two key experiments demonstrated that DNA is the genetic material.
- In the first key experiment (Griffiths, 1928) showed that a virulent strain of Streptococcus pneumoniae. genetically transformed nonvirulent S. pneumoniae. into virulent bacteria.







A. DNA: The Genetic Material

- In a prelude to the second key experiment (Avery, 1944) showed that DNA was the transforming agent through studies of T-even bacteriophage and their treatment with hydrolytic enzymes.
- The second key experiment (Hershey & Chase, 1952) showed that labeled viruses were incubated with host bacteria. Labeled viral DNA entered host cells, producing many label-bearing viruses.









B. The Structure of DNA


 X-ray crystallography showed that the DNA molecule is a helix.






B. The Structure of DNA


- DNA is composed of nucleotides, each containing adenine, cytosine, thymine, or guanine.
- There are equal amounts of adenine and thymine and equal amounts of guanine and cytosine. This is known as Chargaff's Rule (1950, using paper TLC).





B. The Structure of DNA


- Watson and Crick (1953) proposed that DNA is a double-stranded helix with antiparallel strands, and with bases linked by hydrogen bonding.
- Their model accounts for genetic information, mutation, and replication functions of DNA.
















C. DNA Replication


- Semiconservative, conservative, and dispersive models for DNA replication were hypothesized.
- · Each obeyed base-pairing rules.




C. DNA Replication


- Kornberg (1956) demonstrated in vitro that DNA served as its own template during replication.
- Meselson and Stahl's experiment (1957)
 proved replication of DNA to be
 semiconservative. A parent strand is a
 template for synthesis of a new strand. Two
 replicated DNA helices contain one parent
 strand and one synthesized strand each.









- DNA polymerase catalyzes nucleotides from the 5' to the 3' end.
- Nucleotides are added by complementary base pairing with the template strand.
- The substrates, deoxyribonucleoside triphosphates, are hydrolyzed as added, releasing energy for DNA synthesis.





- News Flash: The DNA replication complex is in a fixed location and DNA is threaded through it for replication.
- · Old idea was via moving replication forks.






- Prokaryotes have a single origin of replication; eukaryotes have many (10² to 10³).
- Replication for each proceeds in both directions from an origin of replication.







- Many proteins assist in DNA replication. DNA helicases unwind the double helix, the template strands are stabilized by single-stranded binding proteins.
- An RNA primase catalyzes the synthesis of short RNA primers, and to which nucleotides are added.





- DNA polymerase III action causes the leading strand to grow in the 5'-to-3' direction until replication of that section of DNA is complete.
- RNA primer is degraded and DNA is replaced by DNA polymerase I.


- On the lagging strand, growing in the other direction, DNA is made in the 5'-to-3' direction but synthesis is discontinuous: DNA is added as short Okazaki fragments to primers, then DNA polymerase III skips past the 5' end to make the next fragment.
- DNA polymerase I and Ligase are required to make lagging strand "continuous".













E. DNA Proofreading and Repair


- There is about about one error in 10⁶ nucleotides bases added in DNA replication, repaired by: proofreading, mismatch repair, and excision repair.
- DNA repair mechanisms lower the error rate to about one base in 109.








E. DNA Proofreading and Repair

 Although energetically costly and somewhat redundant, DNA repair is crucial to the survival of the cell.











E. DNA Proofreading and Repair

- Some moderately repetitive DNA sequences, such as telomeric DNA is found at the ends of chromosomes. Some may be lost during each DNA replication, leading to chromosome instability and cell death.
- Telomerase catalyzes the restoration of lost telomeric DNA.
- Most somatic cells lack telomerase and thus have limited life spans.







F. Practical Applications of DNA Replication

- The principles of DNA replication can be used to determine the nucleotide sequence of DNA.
- The polymerase chain reaction technique uses DNA polymerases to repeatedly replicate DNA in the test tube.

