

# 4


## MANIPULATION OF PURIFIED DNA

Once pure samples of DNA have been prepared, the next step in a gene cloning experiment is construction of the recombinant DNA molecule (Figure 1.1). To produce this recombinant molecule, the vector, as well as the DNA to be cloned, must be cut at specific points and then joined together in a controlled manner. Cutting and joining are two examples of DNA manipulative techniques, a wide variety of which have been developed over the last few years. As well as being cut and joined, DNA molecules can be shortened, lengthened, copied into RNA or into new DNA molecules, and modified by the addition or removal of specific chemical groups. These manipulations, all of which can be carried out in the test-tube, provide the foundation not only for gene cloning, but also for basic studies into DNA biochemistry, gene structure, and the control of gene expression.

Almost all DNA manipulative techniques make use of purified enzymes. Within the cell these enzymes participate in essential processes such as DNA replication and transcription, breakdown of unwanted or foreign DNA (for example, invading virus DNA), repair of mutated DNA, and **recombination** between different DNA molecules. After purification from cell extracts, many of these enzymes can be persuaded to carry out their natural reactions, or something closely related to them, under artificial conditions. Although these enzymatic reactions are often quite straightforward, most are absolutely impossible to perform by standard chemical methods. Purified enzymes are therefore crucial to genetic engineering and an important industry has sprung up around their preparation, characterization and marketing. Commercial suppliers of high-purity enzymes provide an essential service to the molecular biologist.

The cutting and joining manipulations that underlie gene cloning are carried out by enzymes called restriction endonucleases (for cutting) and ligases (for joining). Most of this chapter will be concerned with the ways in which these two types of enzyme

are used. Firstly, however, we must consider the whole range of DNA manipulative enzymes, to see exactly what types of reaction can be performed. Many of these enzymes will be mentioned in later chapters when procedures that make use of them will be described.



These enzymes can be grouped into five broad classes depending on the type of reaction that they catalyse.

1. **Nucleases** are enzymes that cut, shorten or degrade nucleic acid molecules.
2. **Ligases** join nucleic acid molecules together.
3. **Polymerases** make copies of molecules.
4. **Modifying enzymes** remove or add chemical groups.
5. **Topoisomerases** introduce or remove supercoils from covalently closed-circular DNA.

Before considering in detail each of these classes of enzyme, two points should be made. The first is that, although most enzymes can be assigned to a particular class, a few display multiple activities that span two or more classes. Most importantly, many polymerases combine their ability to make new DNA molecules with an associated DNA degradative (= nuclease) activity.

Secondly it should be appreciated that, as well as the DNA manipulative enzymes, many similar enzymes able to act on RNA are known. The ribonuclease used to remove contaminating RNA from DNA preparations (p. 33) is an example of such an enzyme. Although some RNA manipulative enzymes have applications in gene cloning and will be mentioned in later chapters, we will in general restrict our thoughts to those enzymes that act on DNA.


#### 4.1.1 Nucleases

Nucleases degrade DNA molecules by breaking the phosphodiester bonds that link one nucleotide to the next in a DNA strand. There are two distinct kinds of nuclease (Figure 4.1).

1. **Exonucleases**, which remove nucleotides one at a time from the end of a DNA molecule.
2. **Endonucleases**, which are able to break internal phosphodiester bonds within a DNA molecule.

The main distinction between different exonucleases lies in the number of strands that are degraded when a double-stranded molecule is attacked. The enzyme called Bal31 (purified from the


**Figure 4.1** The reactions catalysed by the two different kinds of nuclease. (a) An exonuclease, which removes nucleotides from the end of a DNA molecule. (b) An endonuclease, which breaks internal phosphodiester bonds.



bacterium *Alteromonas espejiana*) is an example of an exonuclease that removes nucleotides from both strands of a double-stranded molecule (Figure 4.2(a)). The greater the length of time that Bal31 is allowed to act on a group of DNA molecules, the shorter the resulting DNA fragments will be. In contrast, enzymes such as *E. coli* exonuclease III degrade just one strand of a double-stranded molecule, leaving single-stranded DNA as the product (Figure 4.2(b)).


The same criterion can be used to classify endonucleases. S1 endonuclease (from the fungus *Aspergillus oryzae*) will cleave only single strands (Figure 4.3(a)), whereas deoxyribonuclease I (DNase

D), which is prepared from cow pancreas, cuts both single- and double-stranded molecules (Figure 4.3(b)). DNase I is non-specific in that it attacks DNA at any internal phosphodiester bond; the end result of prolonged DNase I action is therefore a mixture of mononucleotides and very short oligonucleotides. On the other hand, the special group of enzymes called restriction endonucleases cleave double-stranded DNA only at a limited number of specific recognition sites (Figure 4.3(c)). These important enzymes are described in detail on p. 60.




**Figure 4.2** The reactions catalysed by different types of exonuclease. (a) Bal31, which removes nucleotides from both strands of a double-stranded molecule. (b) Exonuclease III, which removes nucleotides only from the 3'-terminus (see p. 81 for a description of the differences between the 3'- and 5'-termini of a polynucleotide).

**Figure 4.3** The reactions catalysed by different types of endonuclease. (a) S1 nuclease, which cleaves only single-stranded DNA, including single-stranded nicks in mainly double-stranded molecules. (b) DNase I, which cleaves both single- and double-stranded DNA. (c) A restriction endonuclease, which cleaves double-stranded DNA, but only at a limited number of sites.



#### 4.1.2 Ligases

In the cell the function of DNA ligase is to repair single-stranded breaks ('discontinuities') that arise in double-stranded DNA molecules during, for example, DNA replication. DNA ligases from most organisms will also join together two individual fragments of



**Figure 4.4** The two reactions catalysed by DNA ligase. (a) Repair of a discontinuity – a missing phosphodiester bond in one stand of a double-stranded molecule. (b) Joining two molecules together.


double-stranded DNA (Figure 4.4). The role of these enzymes in construction of recombinant DNA molecules is described on p. 77.

### 4.1.3 Polymerases

DNA polymerases are enzymes that synthesize a new strand of DNA complementary to an existing DNA or RNA **template** (Figure 4.5(a)). Most polymerases can function only if the template possesses a double-stranded region which acts as a **primer** for initiation of polymerization.

Three types of DNA polymerase are used routinely in genetic engineering. The first is DNA polymerase I, which is prepared usually from *E. coli*. This enzyme attaches to a short single-stranded region (or 'nick') in a mainly double-stranded DNA molecule, and then synthesizes a completely new strand, degrading the existing strand as it proceeds (Figure 4.5(b)). DNA polymerase I is therefore an example of an enzyme with a dual activity – DNA polymerization and DNA degradation.

In fact the polymerase and nuclease activities of DNA polymerase I are controlled by different parts of the enzyme molecule. The nuclease activity is contained in the first 323 amino acids of the polypeptide, so removal of this segment leaves a modified enzyme that retains the polymerase function but is unable to degrade DNA. This modified enzyme, called the **Klenow fragment**, can still synthesize a complementary DNA strand on a single-stranded template,



**Figure 4.5** The reactions catalysed by DNA polymerases. (a) The basic reaction: a new DNA strand is synthesized in the 5' to 3' direction. (b) DNA polymerase I, which initially fills in nicks but then continues to synthesize a new strand, degrading the existing one as it proceeds. (c) The Klenow fragment, which only fills in nicks. (d) Reverse transcriptase, which uses a template of RNA.


but as it has no nuclease activity it cannot continue the synthesis once the nick is filled in (Figure 4.5(c)). Several other enzymes – natural polymerases and modified versions – are able to perform the same function. The major application of the Klenow fragment and these related polymerases is in DNA sequencing (p. 192).

The third type of DNA polymerase that is important in genetic engineering is **reverse transcriptase**, an enzyme involved in the replication of several kinds of virus. Reverse transcriptase is unique in that it uses as a template not DNA but RNA (Figure 4.5(d)). The ability of this enzyme to synthesize a DNA strand complementary to an RNA template is central to the technique called cDNA cloning (p. 167).

### 4.1.4 DNA modifying enzymes

There are numerous enzymes that modify DNA molecules by addition or removal of specific chemical groups. The most important are as follows.

1. **Alkaline phosphatase** (from *E. coli* or calf intestinal tissue) which removes the phosphate group present at the 5'-terminus of a DNA molecule (Figure 4.6(a)).
2. **Polynucleotide kinase** (from *E. coli* infected with T4 phage) which has the reverse effect to alkaline phosphatase, adding phosphate groups on to free 5'-termini (Figure 4.6(b)).
3. **Terminal deoxynucleotidyl transferase** (from calf thymus tissue) which adds one or more deoxynucleotides on to the 3'-terminus of a DNA molecule (Figure 4.6(c)).



**Figure 4.6** The reactions catalysed by DNA-modifying enzymes. (a) Alkaline phosphatase, which removes 5'-phosphate groups. (b) Polynucleotide kinase, which attaches 5'-phosphate groups. (c) Terminal deoxynucleotidyl transferase, which attaches nucleotides to the 3'-termini of polynucleotides in either (i) single-stranded or (ii) double-stranded molecules.

### 4.1.5 Topoisomerases

The final class of DNA manipulative enzymes are the topoisomerases, which are able to change the conformation of covalently closed-circular DNA (e.g. plasmid DNA molecules) by introducing




or removing supercoils (p. 39). Although important in the study of DNA replication, topoisomerases have yet to find a real use in genetic engineering.



Gene cloning requires that DNA molecules be cut in a very precise and reproducible fashion. This is illustrated by the way in which the vector is cut during construction of a recombinant DNA molecule (Figure 4.7(a)). Each vector molecule must be cleaved at a single position, to open up the circle so that new DNA can be inserted; a molecule that is cut more than once will be broken into two or more separate fragments and will be of no use as a cloning vehicle. Furthermore, each vector molecule must be cut at exactly the same position on the circle – as will become apparent in later chapters, random cleavage is not satisfactory. It should be clear

**Figure 4.7** The need for very precise cutting manipulations in a gene cloning experiment.



that a very special type of nuclease is needed to carry out this manipulation.

Often it will also be necessary to cleave the DNA that is to be cloned (Figure 4.7(b)). There are two reasons for this. Firstly, if the aim is to clone a single gene, which may consist of only 2 or 3 kb of DNA, then that gene will have to be cut out of the large (often greater than 80 kb) DNA molecules produced by skilful use of the preparative techniques described in Chapter 3. Secondly, large DNA molecules may have to be broken down simply to produce fragments small enough to be carried by the vector. Most cloning vectors exhibit a preference for DNA fragments that fall into a particular size range; M13-based vectors, for example, are very inefficient at cloning DNA molecules of more than 3 kb in length.


Purified restriction endonucleases allow the molecular biologist to cut DNA molecules in the precise, reproducible manner required for gene cloning. The discovery of these enzymes, which led to Nobel Prizes for W. Arber, H. Smith and D. Nathans in 1978, was one of the key breakthroughs in the development of genetic engineering.

#### 4.2.1 The discovery and function of restriction endonucleases

The initial observation that led to the eventual discovery of restriction endonucleases was made in the early 1950s, when it was shown that some strains of bacteria are immune to bacteriophage infection, a phenomenon referred to as **host-controlled restriction**.

The mechanism of restriction is not very complicated, even though it took over 20 years to be fully understood. Restriction occurs because the bacterium produces an enzyme that degrades the phage DNA before it has time to replicate and direct synthesis of new phage particles (Figure 4.8(a)). The bacterium's own DNA, the destruction of which would of course be lethal, is protected from attack because it carries additional methyl groups that block the degradative enzyme action (Figure 4.8(b)). These degradative enzymes are called restriction endonucleases and are synthesized by many, perhaps all, species of bacteria; over 1200 different ones have now been characterized. Three different classes of restriction endonuclease are recognized, each distinguished by a slightly different mode of action. Types I and III are rather complex and have only a very limited role in genetic engineering. Type II restriction endonucleases, on the other hand, are the cutting enzymes that are so important in gene cloning.

**Figure 4.8** The function of a restriction endonuclease in a bacterial cell. (a) Phage DNA is cleaved, but (b) bacterial DNA is not.



#### 4.2.2 Type II restriction endonucleases cut DNA at specific nucleotide sequences

The central feature of Type II restriction endonucleases (which will be referred to simply as 'restriction endonucleases' from now on) is that each enzyme has a specific recognition sequence at which it cuts a DNA molecule. A particular enzyme will cleave DNA at the recognition sequence and nowhere else. For example, the restriction endonuclease called *PvuI* (isolated from *Proteus vulgaris*) cuts DNA only at the hexanucleotide CGATCG. In contrast, a second enzyme from the same bacterium, called *PvuII*, cuts at a different hexanucleotide, in this case CAGCTG.

Many restriction endonucleases recognize hexanucleotide target sites, but others cut at four, five or even eight nucleotide sequences.

*Sau3A* (from *Staphylococcus aureus* strain 3A) recognizes GATC, and *AluI* (*Arthrobacter luteus*) cuts at AGCT. There are also examples of restriction endonucleases with degenerate recognition sequences, meaning that they cut DNA at any one of a family of related sites. *HinfI* (*Haemophilus influenzae* strain R<sub>f</sub>), for instance, recognizes GANTC, so cuts at GAATC, GATTC, GAGTC and GACTC.

The restriction sequences for some of the most frequently used restriction endonucleases are listed in Table 4.1.

**Table 4.1** The recognition sequences for some of the most frequently used restriction endonucleases

<sup>a</sup> The sequence shown is that of one strand, given in the 5' to 3' direction. Note that almost all recognition sequences are palindromes: when both strands are considered they read the same in each direction, e.g.

```

EcoRI:  5' GAATTC3'
         | | | | |
        3' CTTAAG5'

```


### 4.2.3 Blunt ends and sticky ends

The exact nature of the cut produced by a restriction endonuclease is of considerable importance in the design of a gene cloning experiment. Many restriction endonucleases make a simple double-stranded cut in the middle of the recognition sequence (Figure 4.9(a)), resulting in a **blunt** or flush end. *PvuII* and *AluI* are examples of blunt-end cutters.

However, quite a large number of restriction endonucleases cut DNA in a slightly different way. With these enzymes the two DNA

strands are not cut at exactly the same position. Instead the cleavage is staggered, usually by two or four nucleotides, so that the resulting DNA fragments have short single-stranded overhangs at each end (Figure 4.9(b)). These are called **sticky** or **cohesive ends**, as base pairing between them can stick the DNA molecule back together again (recall that sticky ends were encountered on p. 22 during the description of  $\lambda$  phage replication). One important feature of sticky end enzymes is that restriction endonucleases with different recognition sequences may produce the same sticky ends. *Bam*HI (recognition sequence GGATCC) and *Bgl*III (AGATCT) are examples – both produce GATC sticky ends (Figure 4.9(c)). The

**Figure 4.9** The ends produced by cleavage of DNA with different restriction endonucleases. (a) A blunt end produced by *Acl*I. (b) A sticky end produced by *Eco*RI. (c) The same sticky ends produced by *Bam*HI, *Bgl*III and *Sau*3A.



same sticky end is also produced by *Sau3A*, which recognizes only the tetranucleotide GATC. Fragments of DNA produced by cleavage with either of these enzymes can be joined to each other, as each fragment will carry a complementary sticky end.

#### 4.2.4 The frequency of recognition sequences in a DNA molecule

The number of recognition sequences for a particular restriction endonuclease in a DNA molecule of known length can be calculated mathematically. A tetranucleotide sequence (e.g. GATC) should occur once every  $4^4 = 256$  nucleotides, and a hexanucleotide (e.g. GGATCC) once every  $4^6 = 4096$  nucleotides. These calculations assume that the nucleotides are ordered in a random fashion and that the four different nucleotides are present in equal proportions (i.e. the GC content = 50%). In practice, neither of these assumptions is entirely valid. For example, the  $\lambda$  DNA molecule, at 49 kb, should contain about 12 sites for a restriction endonuclease with a hexanucleotide recognition sequence. In fact these recognition sites occur less frequently (for example, six for *Bgl*III, five for *Bam*HI and only two for *Sal*I), a reflection of the fact that the GC content for  $\lambda$  is rather less than 50% (Figure 4.10(a)).

Furthermore, restriction sites are generally not evenly spaced out along a DNA molecule. If they were then digestion with a particular restriction endonuclease would give fragments of roughly equal sizes. Figure 4.10(b) shows the fragments produced by cutting  $\lambda$  DNA with *Bgl*III, *Bam*HI and *Sal*I. In each case there is a considerable spread of fragment sizes, indicating that in  $\lambda$  DNA the nucleotides are not randomly ordered.


The lesson to be learned from Figure 4.10 is that, although mathematics may give an idea of how many restriction sites to expect in a given DNA molecule, only experimental analysis can provide the true picture. We must therefore move on to consider how restriction endonucleases are used in the laboratory.

#### 4.2.5 Performing a restriction digest in the laboratory

As an example we will consider how to digest a sample of  $\lambda$  DNA (concentration 125  $\mu\text{g}/\text{ml}$ ) with *Bgl*III.

First of all the required amount of DNA must be pipetted into a test-tube. The amount of DNA that will be restricted depends on the nature of the experiment; in this case we shall digest 2  $\mu\text{g}$  of  $\lambda$  DNA, which is contained in 16  $\mu\text{l}$  of the sample (Figure 4.11(a)). Very accurate micropipettes will therefore be needed.

Clearly the other main component in the reaction will be the restriction endonuclease, obtained from a commercial supplier as a pure solution of known concentration. But before adding the



**Figure 4.10** Restriction of the  $\lambda$  DNA molecule. (a) The positions of the recognition sequences for *Bgl*II, *Bam*HI and *Sal*I. (b) The fragments produced by cleavage with each of these restriction endonucleases; the numbers are the fragment sizes in base-pairs.


enzyme, the solution containing the DNA must be adjusted to provide the correct conditions to ensure maximal activity of the enzyme. Most restriction endonucleases function adequately at pH 7.4, but different enzymes vary in their requirements for ionic strength (provided usually by NaCl) and  $Mg^{2+}$  concentration (all Type II restriction endonucleases require  $Mg^{2+}$  in order to function). It is also advisable to add a reducing agent, such as dithiothreitol, which will stabilize the enzyme and prevent its

inactivation. Providing the right conditions for the enzyme is very important – incorrect NaCl or  $Mg^{2+}$  concentrations may not only decrease the activity of the restriction endonuclease, but may also cause changes in the specificity of the enzyme, so that DNA cleavage occurs at additional, non-standard recognition sequences.

The composition of a suitable buffer for *Bgl*III is shown in Table 4.2. This buffer is ten times the working concentration, and is diluted by being added to the reaction mixture. In our example, a suitable final volume for the reaction mixture would be 20  $\mu$ l, so we add 2  $\mu$ l of  $10 \times$  *Bgl*III buffer to the 16  $\mu$ l of DNA already present (Figure 4.11(b)).


The restriction endonuclease can now be added. By convention, 1 unit of enzyme is defined as the quantity needed to cut 1  $\mu$ g of DNA in 1 hour, so we need 2 units of *Bgl*III to cut 2  $\mu$ g of  $\lambda$  DNA. *Bgl*III is frequently obtained at a concentration of 4 units/ $\mu$ l, so 0.5  $\mu$ l will be sufficient to cleave the DNA. The final ingredients in the reaction mixture are therefore 0.5  $\mu$ l *Bgl*III + 1.5  $\mu$ l water, giving a final volume of 20  $\mu$ l (Figure 4.11(c)).

The last factor to consider is incubation temperature. Most



**Figure 4.11** Performing a restriction digest in the laboratory. See the text for details.



**Table 4.2** A 10 × buffer suitable for restriction of DNA with *Bgl*III


restriction endonucleases, including *Bgl*III, work best at 37°C, but a few have different requirements. *Taq*I, for example, is purified from a bacterium called *Thermus aquaticus*, which normally lives at high temperatures in habitats such as hot springs. Restriction digests with *Taq*I must be incubated at 65°C to obtain maximum enzyme activity.

After 1 hour the restriction should be complete (Figure 4.11(d)). If the DNA fragments produced by restriction are to be used in cloning experiments, then the enzyme must somehow be destroyed so that it does not accidentally digest other DNA molecules that may be added at a later stage. There are several ways of 'killing' the enzyme. For many a short incubation at 70°C is sufficient, for others phenol extraction or the addition of EDTA (which binds  $Mg^{2+}$  ions preventing restriction endonuclease action) is used (Figure 4.11(e)).

#### 4.2.6 Analysing the result of restriction endonuclease cleavage

A restriction digest will result in a number of DNA fragments, the sizes of which depend on the exact positions of the recognition sequences for the endonuclease in the original molecule (Figure 4.10). Clearly a way of determining the number and sizes of the fragments is needed if restriction endonucleases are to be of use in gene cloning. Whether or not a DNA molecule is cut at all can be determined fairly easily by testing the viscosity of the solution. Larger DNA molecules result in a more viscous solution than smaller ones, so cleavage is associated with a decrease in viscosity. However, working out the number and sizes of the individual cleavage products is more difficult. In fact, for several years this was one of the most tedious aspects of experiments involving DNA. Eventually the problems were solved in the early 1970s when the technique of gel electrophoresis was developed.

(a) **Separation of molecules by gel electrophoresis** DNA molecules, like proteins and many other biological compounds, carry an electric charge, negative in the case of DNA. Consequently, when DNA molecules are placed in an electric field they migrate towards the positive pole (Figure 4.12(a)). The rate of migration of a molecule depends on two factors, its shape and its charge-to-mass ratio. Unfortunately, most DNA molecules are the same shape and all have very similar charge-to-mass ratios. Fragments of different sizes cannot therefore be separated by standard electrophoresis.



**Figure 4.12** Standard electrophoresis (a) does not separate DNA fragments of different sizes, whereas gel electrophoresis (b) does.

However, the size of the DNA molecule does become a factor if the electrophoresis is performed in a gel. A gel, which is usually made of agarose, polyacrylamide or a mixture of the two, comprises a complex network of pores, through which the DNA molecules must travel to reach the positive electrode. The smaller the DNA molecule, the faster it can migrate through the gel. **Gel electrophoresis** therefore separates DNA molecules according to their size (Figure 4.12(b)).

In practice the composition of the gel determines the sizes of the DNA molecules that can be separated. A 0.5-cm thick slab of 0.5% agarose, which has relatively large pores, would be used for molecules in the size range 1 to 30 kb, allowing, for example, molecules of 10 and 12 kb to be clearly distinguished. At the other end of the scale, a very thin (0.3-mm) 40% polyacrylamide gel, with extremely small pores, would be used to separate much smaller DNA molecules, in the range 1 to 300 bp, and could distinguish molecules differing in length by just a single nucleotide.

#### (b) Visualizing DNA molecules in a gel


(i) *Staining* The easiest way to see the results of a gel electrophoresis experiment is to stain the gel with a compound that will make the DNA visible. Ethidium bromide, already described on p. 42 as a means of visualizing DNA in CsCl gradients, is also routinely used to stain DNA in agarose and polyacrylamide gels (Figure 4.13). Bands showing the positions of the different size classes of DNA fragment are clearly visible under ultraviolet irradiation after EtBr staining, so long as sufficient DNA is present.

(ii) *Autoradiography of radioactively labelled DNA* The one drawback with EtBr staining is that there is a limit to its sensitivity. If less than about 25 ng of DNA is present per band then it is unlikely that the results will show up with EtBr staining. For small amounts of DNA a much more sensitive detection method is needed.

**Autoradiography** provides an answer. If the DNA is labelled, before electrophoresis, by incorporation of a **radioactive marker** into the individual molecules, then the DNA can be visualized by placing an X-ray-sensitive photographic film over the gel. The radioactive DNA exposes the film, revealing the banding pattern (Figure 4.14).

A DNA molecule is usually labelled by incorporating nucleotides that carry a radioactive isotope of phosphorus,  $^{32}\text{P}$  (Figure 4.15(a)). Several methods are available, the most popular being **nick translation** and **end-filling**.

Nick translation refers to the activity of DNA polymerase I (p. 57). Most purified samples of DNA contain some nicked molecules, however carefully the preparation has been carried out, so




**Figure 4.13** Visualizing DNA bands in an agarose gel by EtBr staining and ultraviolet irradiation.

DNA polymerase I will be able to attach to the DNA and catalyse a strand replacement reaction (Figure 4.5(b)). This reaction requires a supply of nucleotides: if one of these is radioactively labelled, then the DNA molecule will itself become labelled (Figure 4.15(b)).

Nick translation can be used to label any DNA molecule but may under some circumstances also cause DNA cleavage. End-filling is a gentler method that rarely causes breakage of the DNA, but unfortunately can only be used to label DNA molecules that have sticky ends. The enzyme used is the Klenow fragment (p. 57), which 'fills in' a sticky end by synthesizing the complementary strand (Figure 4.15(c)). As with nick translation, if the end-filling reaction is carried out in the presence of labelled nucleotides, then the DNA itself becomes labelled.

Both nick translation and end-filling can label DNA to such an extent that very small quantities can be detected in gels by autoradiography. As little as 2 ng of DNA per band can be visualized under ideal conditions.

**Figure 4.14** The use of autoradiography to visualize radioactively labelled DNA in an agarose gel.



#### 4.2.7 Estimation of the sizes of DNA molecules


Gel electrophoresis will separate different-sized DNA molecules, with the smallest molecules travelling the greatest distance towards the positive electrode. If several DNA fragments of varying sizes are present (the result of a successful restriction digest, for example), then a series of bands will appear in the gel. How can the sizes of these fragments be determined?

The most accurate method is to make use of the mathematical relationship that links migration rate to molecular weight. The relevant formula is

$$D = a - b(\log M)$$


where  $D$  is the distance moved,  $M$  is the molecular weight, and  $a$  and  $b$  are constants that depend on the electrophoresis conditions.

However, a much simpler though less precise way of estimating DNA fragment sizes is generally used. A standard restriction digest, comprising fragments of known size, is usually included in each electrophoresis gel that is run. Restriction digests of  $\lambda$  DNA are often used in this way as size markers. For example, *Hind*III cleaves  $\lambda$  DNA into eight fragments, ranging in size from 125 bp for the smallest to over 23 kb for the largest. As the sizes of the fragments in this digest are known, the fragment sizes in the experimental digest can be estimated by comparing the positions of the bands in the two tracks (Figure 4.16). Although not precise, this method can be performed with as little as a 5% error, which is quite satisfactory for most purposes.



**Figure 4.15** Radioactive labelling. (a) The structure of [ $\alpha$ - $^{32}\text{P}$ ]dATP. (b) Labelling DNA by nick translation. (c) Labelling DNA by end-filling.

**Figure 4.16** Estimation of the sizes of DNA fragments in an agarose gel. (a) A rough estimate of fragment size can be obtained by eye. (b) A more accurate measurement of fragment size is gained by using the mobilities of the *Hind*III-fragments to construct a calibration curve; the sizes of the unknown fragments can then be determined from the distances they have migrated.




#### 4.2.8 Mapping the positions of different restriction sites on a DNA molecule

So far we have considered how to determine the number and sizes of the DNA fragments produced by restriction endonuclease cleavage. The next step in **restriction analysis** is to construct a map showing the relative positions on the DNA molecule of the recognition sequences for a number of different enzymes. Only when a **restriction map** is available can the correct restriction endonucleases be selected for the particular cutting manipulation that is required (Figure 4.17).

To construct a restriction map, a series of restriction digests must be performed. Firstly, the number and sizes of the fragments produced by each restriction endonuclease must be determined by gel electrophoresis, followed by comparison with size markers (Figure

4.18). This information must then be supplemented by a series of **double digestions**, in which the DNA is cut by two restriction endonucleases at once. It may be possible to perform a double digestion in one step, if both enzymes have similar requirements for pH,  $Mg^{2+}$  concentration, etc. Alternatively, the two digestions may have to be carried out one after the other, adjusting the reaction mixture after the first digestion to provide a different set of conditions for the second enzyme.

Comparing the results of single and double digests will allow many, if not all, of the restriction sites to be mapped (Figure 4.18). Ambiguities can usually be resolved by **partial digestion**, carried out under conditions that result in cleavage of only a limited number of the restriction sites on any DNA molecule. Partial digestion is achieved normally either by a short incubation period, so the enzyme does not have time to cut all the restriction sites, or by incubation at a low temperature (e.g.  $4^{\circ}C$  rather than  $37^{\circ}C$ ), which limits the activity of the enzyme. The result of a partial digestion is a complex pattern of bands in an electrophoresis gel. As well as the standard fragments, produced by total digestion, additional sizes are seen. These are molecules that comprise two adjacent restriction fragments, separated by a site that has



**Figure 4.17** Using a restriction map to work out which restriction endonucleases should be used to obtain DNA fragments containing individual genes.



not been cleaved. Their sizes indicate which restriction fragments in the complete digest are next to one another in the uncut molecule (Figure 4.18).


**Figure 4.18** Restriction mapping. This example shows how the positions of the *Xba*I, *Xho*I and *Kpn*I sites on the  $\lambda$  DNA molecule can be determined.

#### Single and double digestions

Enzyme	Number of fragments	Sizes (kb)
<i>Xba</i> I	2	24.0 24.5
<i>Xho</i> I	2	15.0 33.5
<i>Kpn</i> I	3	1.5 17.0 30.0
<i>Xba</i> I + <i>Xho</i> I	3	9.0 15.0 24.5
<i>Xba</i> I + <i>Kpn</i> I	4	1.5 6.0 17.0 24.0

#### Conclusions:

- As  $\lambda$  DNA is linear, the number of restriction sites for each enzyme is *Xba*I 1, *Xho*I 1, *Kpn*I 2.
- The *Xba*I and *Xho*I sites can be mapped:



The only possibility is:  $\overline{\text{15.0} \text{ } \overset{\text{XhoI}}{\text{9.0}} \text{ } \overset{\text{XbaI}}{\text{24.5}}}$

- All the *Kpn*I sites fall in the 24.5 kb *Xba*I fragment, as the 24.0 kb fragment is intact after *Xba*I - *Kpn*I double digestion. The order of the *Kpn*I fragments can be determined only by partial digestion.

#### Partial digestion

Enzyme	Fragment sizes (kb)
<i>Kpn</i> I - limiting conditions	1.5, 17.0, 18.5, 30.0, 31.5, 48.5


#### Conclusions:

- 48.5 kb fragment = uncut  $\lambda$ .  
 1.5, 17.0 and 30.0 kb fragments are products of complete digestion.  
 18.5 and 31.5 kb fragments are products of partial digestion.

The *Kpn*I map must be:  $\overline{\text{30.0} \text{ } \overset{\text{KpnI}}{\text{1.5}} \text{ } \text{17.0}}$

Therefore the complete map is:  $\overline{\text{15.0} \text{ } \overset{\text{XhoI}}{\text{9.0}} \text{ } \overset{\text{XbaI}}{\text{6.0}} \text{ } \overset{\text{KpnI}}{\text{1.5}} \text{ } \text{17.0}}$

The final step in construction of a recombinant DNA molecule is the joining together of the vector molecule and the DNA to be cloned (Figure 4.19). This process is referred to as ligation, and the enzyme that catalyses the reaction is called DNA ligase.




**Figure 4.19** Ligation: the final step in construction of a recombinant DNA molecule.

#### 4.3.1 The mode of action of DNA ligase

All living cells produce DNA ligases, but the enzyme used in genetic engineering is usually purified from *E. coli* bacteria that have been infected with T4 phage. Within the cell the enzyme carries out the very important function of repairing any discontinuities that may arise in one of the strands of a double-stranded molecule (Figure 4.4(a)). A discontinuity is quite simply a position where a phosphodiester bond between adjacent nucleotides is missing (contrast this with a 'nick', where one or more nucleotides are absent). Although discontinuities may arise by chance breakage of the cell's DNA molecules, they are also a natural result of processes such as DNA replication and recombination. Ligases therefore play several vital roles in the cell.

In the test-tube, purified DNA ligases, as well as repairing single-strand discontinuities, will also join together individual DNA molecules or the two ends of the same molecule. The chemical reaction involved in ligating two molecules is exactly the same as discontinuity repair, except that two phosphodiester bonds must be made, one for each stand (Figure 4.20(a)).

**Figure 4.20** The different joining reactions catalysed by DNA ligase. (a) Ligation of blunt-ended molecules. (b) Ligation of sticky-ended molecules.



### 4.3.2 Sticky ends increase the efficiency of ligation


The ligation reaction in Figure 4.20(a) shows two blunt-ended fragments being joined together. Although this reaction can be carried out in the test-tube, it is not very efficient. This is because the ligase is unable to 'catch hold' of the molecule to be ligated, and has to wait for chance associations to bring the ends together. If possible, blunt-end ligation should be performed at high DNA concentrations, to increase the chances of the ends of the molecules coming together in the correct way.

In contrast, ligation of complementary sticky ends is much more efficient. This is because compatible sticky ends can base-pair with one another by hydrogen bonding (Figure 4.20(b)) forming a relatively stable structure for the enzyme to work on. If the phosphodiester bonds are not synthesized fairly quickly, then the sticky ends will fall apart again. These transient, base-paired structures do, however, increase the efficiency of ligation by increasing the length of time the ends are in contact with one another.

### 4.3.3 Putting sticky ends on to a blunt-ended molecule

For the reasons detailed in the preceding section, compatible sticky ends are desirable on the DNA molecules to be ligated together in a gene cloning experiment. Often these sticky ends can be provided by digesting both the vector and the DNA to be cloned with the same restriction endonuclease, or with different enzymes that produce the same sticky end. However, it is not always possible to do this. A common situation is where the vector molecule has sticky ends, but the DNA fragments to be cloned are blunt-ended. Under these circumstances one of three methods can be used to put the correct sticky ends on to the DNA fragments.

(a) **Linkers** The first of these methods involves the use of **linkers**. These are short pieces of double-stranded DNA, of known nucleotide sequence, that are synthesized in the test-tube. A typical linker is shown in Figure 4.21(a). It is blunt-ended, but contains a restriction site, *Bam*HI in the example shown. DNA ligase will attach linkers to the ends of larger blunt-ended DNA molecules. Although a blunt-end ligation, this particular reaction can be



**Figure 4.21** Linkers and their use.  
(a) The structure of a typical linker.  
(b) The attachment of linkers to a blunt-ended molecule.


performed very efficiently because synthetic oligonucleotides, such as linkers, can be made in very large amounts and added into the ligation mixture at a high concentration.


More than one linker will attach to each end of the DNA molecule, producing the chain structure shown in Figure 4.21(b). However, digestion with *Bam*HI cleaves the chains at the recognition sequences, producing a large number of cleaved linkers and the original DNA fragment, now carrying *Bam*HI sticky ends. This modified fragment is ready for ligation into a cloning vector restricted with *Bam*HI.

**(b) Adaptors** There is one potential drawback with the use of linkers. Consider what would happen if the blunt-ended molecule shown in Figure 4.21(b) contained one or more *Bam*HI recognition sequences. If this was the case then the restriction step needed to cleave the linkers and produce the sticky ends would also cleave the blunt-ended molecule (Figure 4.22). The resulting fragments will have the correct sticky ends, but that is no consolation if the gene contained in the blunt-ended fragment has now been broken into pieces.

The second method of attaching sticky ends to a blunt-ended molecule is designed to avoid this problem. **Adaptors**, like linkers, are short synthetic oligonucleotides. However, unlike linkers, an adaptor is synthesized so that it already has one sticky end (Figure

**Figure 4.22** A possible problem with the use of linkers. Compare this situation with the desired result of *Bam*HI restriction, as shown in Figure 4.21(b).






**Figure 4.23** Adaptors and the potential problem with their use. (a) A typical adaptor. (b) Two adaptors could ligate to one another to produce a molecule similar to a linker, so that (c) after ligation of adaptors a blunt-ended molecule is still blunt-ended and the restriction step is still needed.

4.23(a)). The idea is of course to ligate the blunt end of the adaptor to the blunt ends of the DNA fragment, to produce a new molecule with sticky ends. This may appear to be a simple method but in practice a new problem arises. The sticky ends of individual adaptor molecules could of course base-pair with each other to form dimers (Figure 4.23(b)), so that the new DNA molecule is still blunt-ended (Figure 4.23(c)). The sticky ends could be recreated by digestion with a restriction endonuclease, but that would defeat the purpose of using adaptors in the first place.

The answer to the problem lies in the precise chemical structure of the ends of the adaptor molecule. Normally, the two ends of a polynucleotide strand are chemically distinct, a fact that will be clear from a careful examination of the polymeric structure (Figure 4.24(a)). One end, referred to as the 5'-terminus, carries a phosphate group (5'-P); the other, the 3'-terminus, has a hydroxyl group (3'-OH). In the double helix the two strands are antiparallel (Figure 4.24(b)), so each end of a double-stranded molecule

**Figure 4.24** The distinction between the 5'- and 3'-termini of a polynucleotide.




consists of one 5'-P terminus and one 3'-OH terminus. Ligation normally takes place between the 5'-P and 3'-OH ends (Figure 4.24(c)).

Adaptor molecules are synthesized so that the blunt end is the same as 'natural' DNA, but the sticky end is different. The 3'-OH terminus of the sticky end is the same as usual, but the 5'-P termi-

nus is modified; it lacks the phosphate group, and is in fact a 5'-OH terminus (Figure 4.25(a)). DNA ligase is unable to form a phosphodiester bridge between 5'-OH and 3'-OH ends. The result is that, although base-pairing is always occurring between the sticky ends of adaptor molecules, the association is never stabilized by ligation (Figure 4.25(b)).

Adaptors can therefore be ligated to a DNA molecule but not to themselves. After the adaptors have been attached the abnormal 5'-OH terminus is converted to the natural 5'-P form by treatment with the enzyme polynucleotide kinase (p. 59), producing a sticky-ended fragment that can be inserted into an appropriate vector.


**(c) Producing sticky ends by homopolymer tailing** The technique of **homopolymer tailing** offers a radically different approach to the production of sticky ends on a blunt-ended DNA molecule. A homopolymer is simply a polymer in which all the subunits are



**Figure 4.25** The use of adaptors. (a) The actual structure of an adaptor, showing the modified 5'-OH terminus. (b) Conversion of blunt ends to sticky ends through the attachment of adaptors.



**Figure 4.26** Homopolymer tailing.  
 (a) Synthesis of a homopolymer tail.  
 (b) Construction of a recombinant DNA molecule from a tailed vector plus tailed insert DNA.  
 (c) Repair of the recombinant DNA molecule.




the same. A DNA strand made up entirely of, say, deoxyguanosine is an example of a homopolymer, and is referred to as poly-deoxyguanosine or poly(dG).

Tailing involves using the enzyme terminal deoxynucleotidyl transferase (p. 59) to add a series of nucleotides on to the 3'-OH termini of a double-stranded DNA molecule. If this reaction is carried out in the presence of just one deoxynucleotide, then a homopolymer tail will be produced (Figure 4.26(a)).

Of course, to be able to ligate together two tailed molecules, the homopolymers must be complementary. Frequently poly(dC) tails are attached to the vector and poly(dG) to the DNA to be cloned. Base-pairing between the two occurs when the DNA molecules are mixed (Figure 4.26(b)).

In practice the poly(dG) and poly(dC) tails are not usually exactly the same length, and base-paired recombinant molecules will have nicks as well as discontinuities (Figure 4.26(c)). Repair is therefore a two-step process, using Klenow polymerase to fill in the nicks followed by DNA ligase to synthesize the final phosphodiester bonds. This repair reaction does not always have to be performed in the test-tube. If the complementary homopolymer tails are longer than about 20 nucleotides, then quite stable base-paired associations will be formed. A recombinant DNA molecule, held together by base-pairing although not completely ligated, will often be stable enough to be introduced into the host cell in the next stage of the cloning experiment (Figure 1.1). Once inside the host, the cell's own DNA polymerase and DNA ligase will repair the recombinant DNA molecule, completing the construction begun in the test-tube.

- 
- Jacobsen, H., Klenow, H. and Overgaard-Hansen, K. (1974) The N-terminal amino-acid sequences of DNA polymerase I from *Escherichia coli* and of the large and small fragments obtained by a limited proteolysis. *European Journal of Biochemistry*, **45**, 623-7 - production of the Klenow fragment of DNA polymerase I.
- Smith, H. O. and Wilcox, K. W. (1970) A restriction enzyme from *Haemophilus influenzae*. *Journal of Molecular Biology*, **51**, 379-91 - one of the first full descriptions of a restriction endonuclease.
- Roberts, R. J. and Macelis, D. (1993) Restriction enzymes and methylases. *Nucleic Acids Research*, **21**, 3125-37 - a comprehensive list of all the known restriction endonucleases and their recognition sequences.
- McDonell, M. W., Simon, M. N. and Studier, F. W. (1977) Analysis of restriction fragments of T7 DNA and determination of molecular weights by electrophoresis in neutral and alkaline gels. *Journal of Molecular Biology*, **110**, 119-46 - an early example of the use of agarose gel electrophoresis in the analysis of restriction fragment sizes.

Lobban, P. and Kaiser, A. D. (1973) Enzymatic end-to-end joining of DNA molecules. *Journal of Molecular Biology*, **79**, 453-71 - ligation.

Rothstein, R. J. *et al.* (1979) Synthetic adaptors for cloning DNA. *Methods in Enzymology*, **68**, 98-109.