



Comparing Prokaryotic and Eukaryotic Cells


Basic unit of living organisms is the cell; the smallest unit capable of life.


“Features” found in all cells:


- Ribosomes
- Cell Membrane
- Genetic Material
- Cytoplasm
- ATP Energy
- External Stimuli
- Regulate Flow
- Reproduce






































TABLE 2.2 Chemical composition of a prokaryotic cell* **Rem: 70-85% Water**

Molecule	Percent of dry weight ^b	Molecules per cell	Different kinds
● Total macromolecules	96	24,610,000	~2500
● Protein	55	2,350,000	~1850
● Polysaccharide	5	4,300	2 ^c
● Lipid	9.1	22,000,000	4 ^d
● Lipopolysaccharide	3.4	1,430,000	1
● DNA	3.1	2.1	1
● RNA	20.5	255,500	~660
Total monomers	3.0		~350
Amino acids and precursors	0.5		~100
Sugars and precursors	2		~50
Nucleotides and precursors	0.5		~200
Inorganic ions	1		18
Total	100%		

^a Data from Neidhardt, F. C., et al. (eds.), 1996. *Escherichia coli* and *Salmonella typhimurium*—Cellular and Molecular Biology, 2nd edition. American Society for Microbiology, Washington, DC.

^b Dry weight of an actively growing cell of *E. coli* is 2.8×10^{-13} g; total weight (70% water) = 9.5×10^{-13} g.

^c Assuming peptidoglycan and glycogen to be the major polysaccharides present.

^d There are several classes of phospholipids, each of which exists in many kinds because of variability in fatty acid composition between species and because of different growth conditions.

Protein ~50%
Lipid ~10% → **Cell Wall 10-20%**
RNA ~20%
DNA ~3-4%

Take Home Message:

Proteins are #1 by weight

Lipids are #1 by number

Peptidoglycan is 1 jumbo molecule

Comparing Prokaryotic and Eukaryotic Cells


Classification of prokaryotic cellular features:
Invariant (or common to all)

⇒ ● **Ribosomes: Sites for protein synthesis – aka the grand translators.**


● **Cell Membranes: The barrier between order and chaos.**

● **Nucleoid Region: Curator of the Information.**

Ribosome structure



(B) Prokaryotic ribosome
(*Escherichia coli*)



(C) Eukaryotic ribosome
(Rat)





Table 7.6 Ribosome structure^a

Property	Prokaryote	Eukaryote
Overall size	70S	80S
Small subunit	30S	40S
Number of proteins	~21	~30
RNA size (number of bases)	16S (1500)	18S (2300)
Large subunit	50S	60S
Number of proteins	~34	~50
RNA size (number of bases)	23S (2900)	28S (4200)
	5S (120)	5.8S (160)
		5S (120)

^a Ribosomes of mitochondria and chloroplasts of eukaryotes are similar to prokaryotic ribosomes (see Section 14.4).

S= Svedberg; a sedimentation coefficient that is NOT ADDITIVE!!!


Protein synthesis




Comparing Prokaryotic and Eukaryotic Cells

Classification of prokaryotic cellular features:
Invariant (or common to all)

- Ribosomes: Sites for protein synthesis – aka the grand translators.
- ➡ ● Cell Membranes: The barrier between order and chaos.
- Nucleoid Region: Curator of the Information.











Table 4.2 Comparative permeability of membranes to various molecules


Substance	Rate of permeability ^a
Water	100
Glycerol	0.1
Tryptophan	0.001
Glucose	0.001
Chloride ion (Cl ⁻)	0.000001
Potassium ion (K ⁺)	0.000001
Sodium ion (Na ⁺)	0.0000001


^a Relative scale—permeability with respect to permeability of water given as 100. Permeability of the membrane to water may be affected by aquaporins (see text).













Archaeal cell membrane structure



Comparing Prokaryotic and Eukaryotic Cells


Classification of prokaryotic cellular features:
Invariant (or common to all)


- Ribosomes: Sites for protein synthesis – aka the grand translators.
- Cell Membranes: The barrier between order and chaos.
- ➡ ● Nucleoid Region: Curator of the Information.





DNA strands released from cell
















Comparing Prokaryotic and Eukaryotic Cells


Classification of prokaryotic cellular features: Variant (or NOT common to all)


- Cell Wall (multiple barrier support themes)
- Endospores (heavy-duty life support strategy)
- Bacterial Flagella (appendages for movement)
- Gas Vesicles (buoyancy compensation devices)
- Capsules/Slime Layer (exterior to cell wall)
- Inclusion Bodies (granules for storage)
- Pili (conduit for genetic exchange)







Cell wall structure








Peptidoglycan of a gram-positive bacterium




















Cell envelopes of *Bacteria*





Structure of the lipopolysaccharide of gram-negative *Bacteria*





The gram-negative cell wall













Morphology of the bacterial endospore
 (a) Terminal (b) Subterminal (c) Central










Table 4.3 Differences between endospores and vegetative cells


Characteristic	Vegetative cell	Endospore
Structure	Typical gram-positive cell; a few gram-negative cells	Thick spore cortex Spore coat Exosporium
Microscopic appearance	Nonrefractile	Refractile
Calcium content	Low	High
Dipicolinic acid	Absent	Present
Enzymatic activity	High	Low
Metabolism (O ₂ uptake)	High	Low or absent
Macromolecular synthesis	Present	Absent
mRNA	Present	Low or absent
Heat resistance	Low	High
Radiation resistance	Low	High
Resistance to chemicals (for example, H ₂ O ₂) and acids	Low	High
Stainability by dyes	Stainable	Stainable only with special methods
Action of lysozyme	Sensitive	Resistant
Water content	High, 80-90%	Low, 10-25% in core
Small acid-soluble proteins (product of <i>sfp</i> genes)	Absent	Present
Cytoplasmic pH	About pH 7	About pH 5.5-6.0 (in core)


Characteristics of Endospore: Take Home Message


- The endospore is a highly resistant differentiated bacterial cell produced by certain gram-positive *Bacteria*.
- Endospore formation leads to a highly dehydrated structure that contains essential macromolecules and a variety of substances such as calcium dipicolinate and small acid-soluble proteins, absent from vegetative cells.
- Endospores can remain dormant indefinitely but germinate quickly when the appropriate trigger is applied.















Flagellar Motility: Take Home Message

- Motility in most microorganisms is due to flagella.
- In prokaryotes the flagellum is a complex structure made of several proteins, most of which are anchored in the cell wall and cytoplasmic membrane.
- The flagellum filament, which is made of a single kind of protein, rotates at the expense of the proton motive force, which drives the flagellar motor.

Gliding Motility: Mechanism??






Gas Vesicles
(a) *Anabaena flos-aquae*
(b) *Microcystis* sp.


The Hammer, Cork, and Bottle Experiment
(Before) (After)



Model of how the two proteins that make up the gas vesicle, *GvpA* and *GvpC*, interact to form a watertight but gas-permeable structure.



Bacterial Capsules: (a) *Acinetobacter* sp. (b) *Rhizobium trifolii*



negative stain

