






More on Phototrophic Potential


























Bacteriochlorophyll Structures

Pigment / Absorption maxima (in vivo)	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇
Bchl a (purple bacteria)/ 805, 830-890	$\begin{array}{c} \text{C}-\text{CH}_3 \\ \\ \text{O} \end{array}$	$-\text{CH}_3^{\text{d}}$	$-\text{CH}_2-\text{CH}_3$	$-\text{CH}_3$	$-\text{C}(=\text{O})-\text{CH}_3$	P/G ^e -H	
Bchl b (purple bacteria)/ 835-850, 1020-1040	$\begin{array}{c} \text{C}-\text{CH}_3 \\ \\ \text{O} \end{array}$	$-\text{CH}_3^{\text{c}}$	$\begin{array}{c} \text{C}-\text{CH}_3 \\ \\ \text{H} \end{array}$	$-\text{CH}_3$	$\begin{array}{c} \text{C}-\text{O}-\text{CH}_3 \\ \\ \text{O} \end{array}$	P	-H
Bchl c (green sulfur bacteria)/745-755	$\begin{array}{c} \text{H} \\ \\ \text{C}-\text{CH}_3 \\ \\ \text{OH} \end{array}$	$-\text{CH}_3$	$-\text{C}_2\text{H}_5^{\text{d}}$	$-\text{C}_2\text{H}_5$	$-\text{C}_2\text{H}_5$	-H	F -CH ₃
Bchl c₂ (green nonsulfur bacteria)/740	$\begin{array}{c} \text{H} \\ \\ \text{C}-\text{CH}_3 \\ \\ \text{OH} \end{array}$	$-\text{CH}_3$	$-\text{C}_2\text{H}_5$	$-\text{CH}_3$	-H	S	-CH ₃
Bchl d (green sulfur bacteria)/705-740	$\begin{array}{c} \text{H} \\ \\ \text{C}-\text{CH}_3 \\ \\ \text{OH} \end{array}$	$-\text{CH}_3$	$-\text{C}_2\text{H}_5$	$-\text{C}_2\text{H}_5$	$-\text{C}_2\text{H}_5$	-H	F -H
Bchl e (green sulfur bacteria)/719-726	$\begin{array}{c} \text{H} \\ \\ \text{C}-\text{CH}_3 \\ \\ \text{OH} \end{array}$	$-\text{C}_2\text{H}_5$	$-\text{C}-\text{H}$	$-\text{C}_2\text{H}_5$	$-\text{C}_2\text{H}_5$	-H	F -CH ₃
Bchl g (halobacteria)/ 670, 788	$\begin{array}{c} \text{H} \\ \\ \text{C}-\text{CH}_2 \\ \\ \text{O} \end{array}$	$-\text{CH}_3^{\text{d}}$	$-\text{C}_2\text{H}_5$	$-\text{CH}_3$	$-\text{C}(=\text{O})-\text{CH}_3$	F	-H

^aP, Phytol ester (C₂₀H₃₉O—); F, farnesyl ester (C₁₅H₂₅O—); Gg, geranylgeranyl ester (C₂₄H₄₁O—); S, stearyl alcohol (C₁₈H₃₇O—).

^bNo double bond between C₃ and C₄; additional H atoms are in positions C₃ and C₄.

^cNo double bond between C₃ and C₄; an additional H atom is in position C₃.

^dBacteriochlorophylls c, d, and e consist of isomeric mixtures with the different substituents on R₃ as shown.







Table 9.1 Some general properties of the various photosynthetic bacteria


	Nonsulfur Purple Bacteria	Purple Sulfur Bacteria	Green Sulfur Bacteria	Cyano-bacteria	Helio-bacteria
Source of reducing power (e^-)	H_2 , reduced organic	H_2S	H_2S	H_2O	Lactate, organic
Oxidized product	Oxidized organic	SO_4^{2-}	SO_4^{2-}	O_2	Oxidized organic
Source of carbon	CO_2 or organic	CO_2	CO_2	CO_2	Lactate pyruvate
Heterotrophic growth	Common	Limited ^d	Limited ^d	Limited ^d	Required

^dGenerally limited to assimilation of low molecular weight organics during autotrophic growth.
















Table 9.2 The bacteriochlorophyll present in photosynthetic bacteria and primary acceptors involved in energy conserving reactions

	Electron Donor	Electron Acceptor
Purple nonsulfur bacteria	Bacteriochlorophyll <i>a</i> and <i>b</i>	Bacteriopheophytin <i>a</i> , Q_A , and Q_B
Green sulfur bacteria	Bacteriochlorophyll <i>c</i> , <i>d</i> , and <i>e</i>	Bacteriopheophytin <i>a</i> and FeS-protein
Cyanobacteria photosystem I	Chlorophyll <i>a</i>	Chlorophyll <i>a</i> and FeS-protein
Cyanobacteria photosystem II	Chlorophyll <i>a</i>	Pheophytin <i>a</i> , Q_A , Q_B , and plastoquinones
<i>Heliobacteria</i>	Bacteriochlorophyll <i>g</i>	Bacteriochlorophyll <i>c</i> and FeS-protein












Absorption Spectra for Cyanobacteria







Reaction center of purple nonsulfur bacteria




Sulfur granules in purple sulfur bacteria
e.g., *Chromatium*




Sulfur granules in green sulfur bacteria
e.g., *Chlorobium*




Electron flow in phototrophs



Reverse TCA in GSBs




Hydroxypropionate in GNBs






A light-driven proton pump of halophilic archaea



Archaeal ~~XXXX~~ rhodopsin molecule has seven helical regions.

Light-driven proton pump of halophilic archaea



Archaeal rhodopsin: retinal structure



Proteorhodopsin in marine *Bacteria* and *Archaea*

