The Process of Growth

- Metabolism required for growth, both anabolic and catabolic. ~2000 reactions!
- Usual Definition: **Increase in cell numbers**
  Other definitions possible – spores, UMC's, respiration, viable but nonculturable, morphology changes (life cycle)
- Divide via Binary Fission: 3 mechanisms involved!
  Cell Elongation – cell wall
  DNA Replication – rate limiting step
  Cell Division – septum formation





Figure 1

Effect of nutrition-imposed growth rate on the composition of *Escherichia coli* B/r. All values are expressed in amounts per cell normalized to values at $\mu = 0.6$ (mass = 1.48×10^{-13} g; protein = 1.00×10^{-13} g; RNA = 2.0×10^{-14} g; DNA = 6.3×10^{-15} g). (Plotted from data in Bremer and Dennis, 1987.)



Figure 2

Effect of growth rate on the cellular proportions of protein, RNA, and DNA. Filled circles refer to results from cultures undergoing balanced growth in batch culture in various media; open circles are from cultures growing in a glucose-limited chemostat. (From Jacobsen, 1974.)

FtsZ ring & cell division

The "divisome"



phase

DNA stain

FtsZ stain

DNA & FtsZ

FtsZ similar to Tubulin
MreB similar to Actin







Bactoprenol, the lipid carrier of peptidoglycan bulding blocks.



Penicillin blocks this reaction





Figure 10

Chromosome structure and equivalent DNA content of the average cell in culture of *E. coli* B/r growing at various rates. The numbers represent genome equivalents. (From Bremer and Dennis, 1987.)



Figure 14

Electron micrograph of a mixture of cells of *E. coli* B/r grown at different rates. The large cells grew with a doubling time of 22 minutes, the small ones with a doubling time of 72 minutes. (From Nanninga and Woldringh, 1985.)

The Process of Growth

- Growth Rate: Time it takes to reproduce
  $t_{\frac{1}{2}} = \ln 2/\mu = 0.693/\mu = g$
- Phases of Growth in Batch culture
  Lag, Log, Stationary, Death
- Measurement of Growth
  Total cell counts
  Viable cell counts
  Turbidity

The growth rate of a microbial culture

Time (h)	Total number of cells	Time (h)	Total number of cells
0	1	4	256 (2^8)
0.5	2	4.5	512 (2^9)
1	4	5	1,024 (2^{10})
1.5	8	5.5	2,048 (2^{11})
2	16	6	4,096 (2^{12})
2.5	32	.	.
3	64	.	.
3.5	128	10	1,048,576 (2^{19})



How to estimate the generation times of an exponential microbial culture using semi-log plots.





Table 6.1 Approximate generation times for several organisms growing in media optimal for growth

Species	Generation Time
Escherichia coli	20 min
Bacillus subtilis	28 min
Staphylococcus aureus	30 min
Pseudomonas aeruginosa	35 min
Thermus aquaticus	50 min
Thermoproteus tenax	1 hr 40 min
Rhodobacter sphaeroides	2 hr 20 min
Sulfolobus acidocaldarius	4 hr
Thermoleophilum album	6 hr
Thermofilum pendens	10 hr
Mycobacterium tuberculosis	13 hr 20 min

Cryptic Growth ⬇



Typical growth curve for a bacterial population

Total Cell counts using the Petroff-Hausser Counter



Total Cell counts using the Petroff-Hausser Counter



Viable cell count methods

30-300 on standard Petri Dish



Counting the number of viable cells by serial dilution and plate count



(Part 1) Concentration of cells by membrane filtration



(Part 2) Concentration of cells by membrane filtration

Turbidity measurements of microbial growth



Turbidity measurements of microbial growth



The Process of Growth

- Continuous Culture: The wonders of the **chemostat**
  - Steady State
  - Reproducible Physiology
  - Fine control

  Key parameters: Ks, μmax, Yield

  Closed systems vs. Open systems vs. Nature!





D = F/V = μ

Where:
D = dilution rate
F = flow rate
V = volume
μ = growth rate

Rem: At Steady State







Fig. 6.10 *Dependence of growth rate* μ *on the substrate concentration* c_s.

Steady-state relationship between substrate concentration and output of bacterial mass



Table 6.2 Growth yields of anaerobic bacteria utilizing glucose as the energy source

	Mol ATP/Mol Glucose	y_{max} (g of cell/mol Glucose)	y_{ATP} (g of cell/mol ATP)
Lactobacillus delbrueckii[a]	2	21	10.5
Enterococcus faecalis[a]	2	20	10
Zymomonas mobilis[b]	1	9	9

[a]Homolactic fermentation, Embden–Meyerhof pathway (see Chapter 10).
[b]Alcoholic fermentation, Entner–Doudoroff pathway (see Chapter 10).