

## Microbes as Energy Transducers

- The Metabolic Menu
- Metabolic Strategies
- Respiration & Fermentation
- Chemolithotrophy
- Photoautotrophy
- Biogeochemical Cycles
- Metabolism in Primitive Organisms


All major types of nutrition and metabolism evolved among prokaryotes: they are the ultimate biochemists

The prokaryotes exhibit some unique modes of nutrition as well as every type of nutrition found in eukaryotes.

### Major Modes of Nutrition:



Prokaryotes exhibit a great diversity in how they obtain the necessary resources (**energy and carbon**) to synthesize organic compounds.


- Some obtain energy from light (**phototrophs**), while others use chemicals taken from the environment (**chemotrophs**).
- Many can utilize CO<sub>2</sub> as a carbon source (**autotrophs**) and others require at least one organic nutrient as a carbon source (**heterotrophs**).



Depending upon the energy source **AND** the carbon source, prokaryotes have **four** possible nutritional modes:





- 1. Photoautotrophs:** Use light energy to synthesize organic compounds from CO<sub>2</sub> – Includes the cyanobacteria. (Actually all photosynthetic eukaryotes fit in this category.)
- 2. Chemoautotrophs:** Require only CO<sub>2</sub> as a carbon source and obtain energy by oxidizing inorganic compounds. This mode of nutrition is unique only to certain prokaryotes.
- 3. Photoheterotrophs:** Use light to generate ATP from an organic carbon source. This mode of nutrition is unique only to certain prokaryotes.
- 4. Chemoheterotrophs:** Must obtain organic molecules for energy and as a source of carbon. Found in many bacteria as well as most eukaryotes.












**Metabolic Menu For Chemotrophs**

Potential Microbial Metabolic Processes			
e <sup>-</sup> donor	e <sup>-</sup> acceptor	C source	Organisms
<b>Autolithotrophy</b>			
H <sub>2</sub>	O <sub>2</sub>	CO <sub>2</sub>	Hydrogen oxidizers
HS <sup>-</sup> /S <sub>2</sub> O <sub>3</sub> <sup>2-</sup>	O <sub>2</sub>	CO <sub>2</sub>	Sulfur oxidizers
Fe <sup>2+</sup>	O <sub>2</sub>	CO <sub>2</sub>	Iron oxidizers
Mn <sup>2+</sup>	O <sub>2</sub>	CO <sub>2</sub>	Manganese oxidizers
NH <sub>4</sub> <sup>+</sup> /NO <sub>2</sub> <sup>-</sup>	O <sub>2</sub>	CO <sub>2</sub>	Nitrifiers
HS <sup>-</sup> /S <sub>2</sub> O <sub>3</sub> <sup>2-</sup>	NO <sub>3</sub> <sup>-</sup>	CO <sub>2</sub>	Denitrifying S-oxidizers
H <sub>2</sub>	NO <sub>3</sub> <sup>-</sup>	CO <sub>2</sub>	Hydrogen oxidizers
H <sub>2</sub>	S <sup>0</sup> /S <sub>2</sub> O <sub>3</sub> <sup>2-</sup>	CO <sub>2</sub>	Sulfate Reducers (SRBs)
H <sub>2</sub>	CO <sub>2</sub>	CO <sub>2</sub>	Methanogens & Acetogens
<b>Heteroorganotrophy</b>			
Org.C	O <sub>2</sub>	Org.C	Aerobic Heterotrophy
Org.C	NO <sub>3</sub> <sup>-</sup>	Org.C	Denitrifiers
Org.C	S <sup>0</sup> /S <sub>2</sub> O <sub>3</sub> <sup>2-</sup>	Org.C	Sulfate Reducers (SRBs)
Org.C	Org.C	Org.C	Fermenters
<b>Methylotrophy</b>			
CH <sub>4</sub> (C-1)	O <sub>2</sub> /SO <sub>4</sub> <sup>2-</sup>	CH <sub>4</sub> /CO <sub>2</sub> /CO	Methane (C-1) oxidizers












### 7.1 Cellular Locations for Energy Pathways in Eukaryotes and Prokaryotes

EUKARYOTES	PROKARYOTES
<b>External to mitochondrion</b>	<b>In cytoplasm</b>
Glycolysis	Glycolysis
Fermentation	Fermentation
	Citric acid cycle
<b>Inside mitochondrion</b>	<b>On inner face of plasma membrane</b>
Inner membrane	Pyruvate oxidation
Pyruvate oxidation	Respiratory chain
Respiratory chain	
Matrix	
Citric acid cycle	









Overall reaction:  $\text{Pyruvate}^- + 4 \text{NAD}^+ + \text{FAD} \rightarrow 3 \text{CO}_2 + 4 \text{NADH} + \text{FADH}$





(1) Substrate-level phosphorylation  $\text{GDP} + \text{P}_i \rightarrow \text{GTP}$   
 $\text{GTP} + \text{ADP} \rightarrow \text{GDP} + \text{ATP}$  } 15 ATP

(2) Electron transport phosphorylation  $4 \text{NADH} \equiv 12 \text{ATP}$   
 $\text{FADH} \equiv 2 \text{ATP}$

(3) Sum: CAC plus glycolysis  $\rightarrow 38 \text{ATP}$  per glucose

(b)





### Table 9.2 ATP Yield from the Aerobic Oxidation of Glucose by Eucaryotic Cells

<b>Glycolytic Pathway</b>	
Substrate-level phosphorylation (ATP)	2 ATP <sup>a</sup>
Oxidative phosphorylation with 2 NADH	6 ATP
<b>2 Pyruvate to 2 Acetyl-CoA</b>	
Oxidative phosphorylation with 2 NADH	6 ATP
<b>Tricarboxylic Acid Cycle</b>	
Substrate-level phosphorylation (GTP)	2 ATP
Oxidative phosphorylation with 6 NADH	18 ATP
Oxidative phosphorylation with 2 $FADH_2$	4 ATP
<b>Total Aerobic Yield</b>	<b>38 ATP</b>

<sup>a</sup>ATP yields are calculated with an assumed P/O ratio of 3.0 for NADH and 2.0 for  $FADH_2$ .

### Fermentation – Key Features



- (1) Substrate-level phosphorylation is the rule\*.
- (2) Always anaerobic (even when some  $O_2$  might be around).
- (3) No externally supplied terminal electron acceptor.

**Many types.... 2 major themes**

- (1)  $NADH+H^+$  gets oxidized to  $NAD^+$
- (2) Electron acceptor is usually **Pyruvate** or its derivative.



\*Rules are always meant to be broken!

**Pasteur Effect:** ~20X more biomass when aerated




**Figure 9.9 Reoxidation of NADH During Fermentation.** NADH from glycolysis is reoxidized by being used to reduce pyruvate or a pyruvate derivative (X). Either lactate or reduced product Y result.


**Figure 9.10 Some Common Microbial Fermentations.** Only pyruvate fermentations are shown for the sake of simplicity; many other organic molecules can be fermented. Most of these pathways have been simplified by deletion of one or more steps and intermediates. Pyruvate and major end products are shown in color.



**(a) Alcohol fermentation**



**(b) Lactic acid fermentation**



**Fig. 8.3. Methylmalonyl-CoA pathway of propionate formation.**  
Enzymes: (1) lactate dehydrogenase; (2) methylmalonyl-CoA carboxytransferase; (3) malate dehydrogenase; (4) fumarase; (5) fumarate reductase (leading to regeneration of ATP by proton translocation); (6) CoA transferase; (7) methylmalonyl-CoA mutase.

