






# Comparing Bacteria, Archaea and Eucarya


## **Classification of microbial cellular features:**

### **Variant (or NOT common to all)**


- Cell Wall (multiple barrier support themes)
- Endospores (heavy-duty life support strategy)
- Bacterial Flagella (appendages for movement)
- Gas Vesicles (buoyancy compensation devices)
- Capsules/Slime Layer (exterior to cell wall)
- Inclusion Bodies (granules for storage)
- Pili (conduit for genetic exchange)

# Bacterial morphologies






# Famous Walls




# Cell walls of *Bacteria*

**Gram-positive**



(a)

**Gram-negative**




(b)




# Cell wall structure



(A) Gram-positive have single-layer cell wall.



(B) Gram-negative have two-layer cell wall.




# Peptidoglycan




Constant: glycan (NAM and NAG),  $\beta$ -1,4 glycosidic linkages horizontally


Variable: type of peptide bridge, connects sheets vertically



# Peptidoglycan of a gram-positive bacterium









(a) DAP or Diaminopimelic acid

Crossing linking AAs




(b) Lysine



(a) *Escherichia coli*  
(gram-negative)

(b) *Staphylococcus aureus*  
(gram-positive)




(c)


## Overall structure of peptidoglycan

# Cell walls of gram-positive and gram-negative bacteria


(A) Gram-positive peptidoglycan



(B) Gram-negative peptidoglycan




# Summary diagram of the gram-positive cell wall



(b)






## Teichoic acids and the overall structure of the gram-positive cell wall

- repeating units of this structure
- negatively charged, contribute to negative charge of cell surface
- found in wall, membrane, and capsule
- may be covalently attached to membrane lipids

(a)


# Cell envelopes of *Bacteria*

(A) Gram-positive cell envelope




# Cell envelopes of *Bacteria*

(B) Gram-negative cell envelope




# Structure of the lipopolysaccharide of gram-negative *Bacteria*





# LPS chemical structure varies by species



-Hydrophilic, so exclude hydrophobic molecules like antibiotics and bile salts.

-Can be toxic

-Serve as an “epitope” or “antigen”


-Antibodies to intact cells’ O-antigen are VERY strain-specific.

-Embedded in lipid layer.

-“Endotoxin”, causes fever and shock in mammals if released from membrane (when bacteria lyse)

-Free (whole) LPS also triggers host defense by binding a receptor in macrophages

# The gram-negative cell wall






## Cell walls of *Archaea*:


Some *Archaea* & *Bacteria* have a protein jacket outside the membrane called the “paracrystalline surface layer” or S-layer.

The S-layer sometimes serves as **cell wall** for *Archaea*.

It is a selective sieve, allowing the passage of low-molecular-weight substances while excluding large molecules and structures.






...other *Archaea* have polysaccharide cell walls






# Formation of the endospore




Spores remain viable in the environment after long periods of dormancy.

Extreme reports of endospore revival (successful germination from):


1. *Bacillus sphaericus* found in the guts of bees preserved in 40 million year old Dominican amber
2. *Virgibacillus* spp. found in salt crystals in the 250 million year old Salado Formation in New Mexico



(a)



(b)



(c)

Morphology of the bacterial endospore  
(a) Terminal (b) Subterminal (c) Central

**Endospores** are a highly resistant differentiated bacterial cell produced by **certain gram-positive *Bacteria***.

- mostly soil bacteria of phylum Firmicutes
- evolved just once
- most common in *Clostridium*, *Bacillus*
- agents of **survival**
- metabolically inert, highly dehydrated (10-15% water)
- most resistant biological structure known: heat up to 150°C, dryness, UV, strong acids, disinfectants
- can survive 100's (thousands? millions?) of years

**Exospores** are formed by pinching off of tips of filamentous bacteria (and of fungi)

- Streptomyces*, *Myxobacteria*
- agents of **dispersal**









Figure 4.25c Microbiology: An Evolving Science  
 © 2009 W. W. Norton & Company, Inc.

*Bacillus megaterium*




*Bacillus subtilis*






**Figure 3.45 Endospore Structure.** *Bacillus anthracis* endospore ( $\times 151,000$ ). Note the following structures: exosporium, EX; spore coat, SC; cortex, CX; core wall, CW; and the protoplast or core with its nucleoid, N, and ribosomes, CR.



(a)



(b)

(a) Structure of Dipicolinic Acid & (b) crosslinked with  $\text{Ca}^{++}$




**Table 4.3****Differences between endospores and vegetative cells**


<b>Characteristic</b>	<b>Vegetative cell</b>	<b>Endospore</b>
Structure	Typical gram-positive cell; a few gram-negative cells	Thick spore cortex Spore coat Exosporium
Microscopic appearance	Nonrefractile	Refractile
Calcium content	Low	High
● Dipicolinic acid	Absent	Present
Enzymatic activity	High	Low
Metabolism (O <sub>2</sub> uptake)	High	Low or absent
Macromolecular synthesis	Present	Absent
mRNA	Present	Low or absent
Heat resistance	Low	High
Radiation resistance	Low	High
Resistance to chemicals (for example, H <sub>2</sub> O <sub>2</sub> ) and acids	Low	High
Stainability by dyes	Stainable	Stainable only with special methods
Action of lysozyme	Sensitive	Resistant
Water content	High, 80–90%	Low, 10–25% in core
● Small acid-soluble proteins (product of <i>ssp</i> genes)	Absent	Present
Cytoplasmic pH	About pH 7	About pH 5.5–6.0 (in core)

# Characteristics of Endospore: Take Home Message

- The endospore is a highly resistant differentiated bacterial cell produced by certain gram-positive *Bacteria*.
- Endospore formation leads to a highly dehydrated structure that contains essential macromolecules and a variety of substances such as calcium dipicolinate and small acid-soluble proteins, absent from vegetative cells.
- Endospores can remain dormant indefinitely but germinate quickly when the appropriate trigger is applied.




**Bacterial flagella**  
**(A) Polar**  
**(aka monotrichous)**  
**&**  
**(B) Peritrichous**




**Bacterial flagella cont.**

Also:

(C) Amphitrichous (bipolar)  
&  
(D) Lophotrichous (tuft)








## Structure of the bacterial flagellum

Bacteria: 60 cell lengths/second  
 Cheetah: 25 body lengths/second




## Proton Transport-Coupled Rotation of the Flagellum




Model for Flagellar Rotation: Mot proteins (MotA/MotB complex) form two half-channels.

Estimate:  $\sim 1000$  protons/1 turn




**The Ultrastructure of Bacterial Flagella.** Flagellar basal bodies and hooks in (a) gram-negative and (b) gram-positive bacteria.




Flagellar Motility: Relationship of flagellar rotation to bacterial movement.



➔ **Reversible flagella**




➔ **Unidirectional flagella**



**Polar** (both)

Flagellar Motility: Relationship of flagellar rotation to bacterial movement.

# Flagellar Assembly




**Growth of Flagellar Filaments.** Flagellin subunits travel through the flagellar core and attach to the growing tip.


## Flagellar Motility: Take Home Message

- Motility in most microorganisms is due to flagella.
- In bacteria the flagellum is a complex structure made of several proteins, most of which are anchored in the cell wall and cytoplasmic membrane.
- The flagellum filament, which is made of a single kind of protein, rotates at the expense of the proton motive force, which drives the flagellar motor.


# Gliding Motility: Mechanism??







A




B

Gas Vesicles  
(A) *Anabaena flos-aquae*  
(B) *Microcystis* sp.


# The Hammer, Cork, and Bottle Experiment

(Before) (After)







Model of how the two proteins that make up the gas vesicle, *GvpA* and *GvpC*, interact to form a watertight but gas-permeable structure.




Bacterial Capsules: (A) *Acinetobacter* sp. (B) *Rhizobium trifolii*




negative stain






Storage of PHB




Poly-β-hydroxybutyrate




Sulfur globules inside the purple sulfur bacterium *Isochromatium buderi*

Magnetotactic bacteria with  $\text{Fe}_3\text{O}_4$  (magnetite) particles called magnetosomes










**EM of *Salmonella typhi***





"Sex" Pili used in bacterial conjugation of *E. coli* cells

# Carboxysomes of *Thiobacillus neapolitanus*



- Carboxysomes found in autotrophs
- Cytoplasmic inclusion involved in fixation of CO<sub>2</sub> into cellular biomass

## Extensive cell membranes of *Nitrosococcus*:



Complex membrane structures of many chemolithotrophs and photoautotrophs – analogous to thylakoid membranes of chloroplasts.

Close linkage between e- transport (fueled by photon capture or oxidation of inorganic compound) and fixation of CO<sub>2</sub>