

## General Patterns in Evolution



---

---

---

---

---

---

---

---

## Uses of Phylogenetic Analysis

- Allows mapping order of character state changes
- Documents evolutionary trends in development
- Reveals that Homoplasy is common
- Can attempt to equate timing with fossil record events

---

---

---


---

---

---

---

---



---

---

---

---


---

---

---

---

Tracing the path of evolution to *Homo sapiens* from the universal ancestor of all life



1. Origin of eukaryotes: a symbiotic bacterium becomes the mitochondrion.
2. Multicellularity evolves: cell and tissue differentiation.
3. Animals: internal digestive cavity develops.
4. Deuterostomes: embryonic blastopore develops into anus.
5. Chordates: notochord, dorsal nerve cord.
6. Vertebrates: brain develops.
7. Synapsids: legs.
8. Amniotes: amniotic egg, other water conserving features.
9. Mammals: unique jaw joints, milk teeth, mammary milk.
10. Primates: binocular vision, adaptability.
11. Anthropoid apes: loss of tail.
12. Hominins: evolve bipedality.
13. Homo sapiens: spreads from Africa.

---

---

---

---

---

---


---

---

---

---

Phylogenetic Analysis Documents Evolutionary Trends in Development: In fruit flies




---

---

---

---

---

---

---


---

---

---

Phylogenies Reveal that Homoplasy is Common

• **Convergent** and **parallel** evolution - the independent gain of a trait




---

---

---

---

---

---


---

---

---

---

### Convergent Evolution among Placental Mammals and Marsupials




---

---

---

---


---

---

---

---

### Structure of the p115RhoGEF rgRGS domain- $\alpha$ 13/11 chimera complex suggests convergent evolution of a GTPase activator.




---

---

---

---

---


---

---

---

### Parallel evolution: Special case of Convergent evolution

Feeding structures (maxillipeds) from thoracic legs in crustaceans.




---

---

---

---

---


---

---

---

**Phylogenies Reveal that Homoplasy is Common**

- Evolutionary reversal - the loss of a trait




---

---

---

---

---


---

---

---

**Reversal:**

An example of the acquisition and loss of individualization  
Homodonts vs. Heterodonts




---

---

---

---

---

---

---

---



**Reconciling the Fossil Record  
with phylogenetic analysis?**

Can really only work with  
morphology-based cladistics.

These images taken from Heck's  
*Iconographic Encyclopedia* (1851).




---

---

---

---

---



---

---


---

## The Evolution of Traits aka phenotypes

Linear interactions



Non-linear interactions




---

---

---

---

---

---

---

---


---

---

---

---

Polymorphic mimicry in *Papilio dardanus* (The mocker swallowtail): accurate mimics of different species of distasteful butterflies.



Single Locus; ~11 mimicking alleles that are more variable.

---

---

---

---

---

---

---

---

---

---

---

---

Biological traits come about through developmental processes and physiological regulatory mechanisms. Most of these processes are nonlinear. Examples of nonlinear processes are:

- The sensitivity of reaction rate to substrate concentration
- inhibition
- negative feedback
- positive feedback
- cooperativity
- most non-steady state processes
- any process that depends on diffusion

Any mechanism that contains one or more of these processes (and most regulatory mechanisms in biology do) will have a nonlinear relationship between variation in its determinants and variation in the trait affected by the process.

---

---

---

---

---

---

---

---


---

---

---

---

**Rates of Evolution Vary Among Lineages**




---

---

---

---

---

---

---

---

**Rates of Evolution Vary Among Characters**



Evolution of different characters  
at different rates within a lineage:  
**Mosaic Evolution**

Combines concepts of  
**Gradualism vs. Saltation**

---

---

---


---

---

---

---

---



Inferred pairwise nucleotide substitutions among 17 mammal species from seven gene products, as estimated from protein studies, plotted against date of divergence, as estimated from the fossil record. The line is drawn from the origin through the oldest point (marsupial/placental divergence at 125 MYA). The strong linear relationship suggests that **molecular differences between pairs of species are proportional to the time of their separation**, rather than the degree of organismal difference. Therefore, measures of genetic divergence can be used to date the time of divergence for species pairs for which no fossil data are available: genes function as **Molecular Clocks**. (from A. C. Wilson 1976)

---

---

---

---

---

---

---

---

**Change in Form is Often Correlated with Change in Function**




Figure 3.9 The panda's thumb (a) Giant pandas strip the leaves from bamboo by passing the stalk through their hands. (Bill Kamin/Visuals Unlimited) (b) This drawing shows how the panda's "thumb" forms a slot for bamboo stalks to pass through. After Endo et al. 1999.

---

---

---

---

---

---

---

---

---

---

**Heterochrony:  
Changes in the Rate or Timing of Developmental Events**



**Paedomorphosis:** the retention of juvenile features in the reproductive adult.

**Peramorphosis:** 'hyper-adult' features in the reproductive adult.

---

---

---

---

---

---

---

---

---

---

**Heterotopy:  
Changes in the Position in which a Trait is Expressed**

Philodendron switching stem and root positions.




---

---

---

---

---

---

---

---

---

---





$\ln(Y) = Y'$

$a > 1$   
positive allometry

$a = 1$

Positive Allometry:  
Y grows quickly relative to X

$\ln(X) = X'$

Legs grow quickly relative to torso

Are we just baby chimps?  
A tale of heterochrony and allometric growth.

*Homo sapiens*, whose prolonged brain development period and relatively flat face may be reflections of a prolonged juvenile period, relative to that of our closest relatives, the bonobos and chimpanzees (*Pan paniscus* and *P. troglodytes*).

---

---

---

---

---

---

---

---

Allometric differences in the jaws among three closely related families of fishes.

(A) Flying fish

(B) Halibut

(C) Needlefish

---

---

---

---

---

---

---

---

**Adaptive Radiation is Widespread  
aka Divergent Evolution**

Hawaiian Honeycreepers

*Reophaeus carolinensis* Laysan Finch  
*Chamaea fasciata* Akaka  
*Myzomela olivacea* Hawaii Amakihi  
*P. palmeri* Palmyra Amakihi  
*Loxia coccinea* Laysan Amakihi  
*L. vitiensis* Laysan Amakihi  
*Melospiza coccinea* Laysan Amakihi  
*Palmeria alba* Palmyra Amakihi  
*Himatione sanguinea* Hawaii Amakihi  
*Psaltriparus minimus* Laysan Amakihi

---

---

---

---

---

---

---

---







---

---

---

---

---

---

---

---