




Mass Extinctions & Their Consequences



Taxonomic diversity of skeletonized marine animal families during the Phanerozoic



Spindle diagram of family diversification/extinction



The role of extinction in evolution

DAVID M. RAUP

Department of Geophysical Sciences, University of Chicago, Chicago, IL 60637


ABSTRACT The extinction of species is not normally considered an important element of modern-day theory, in contrast to the opposite phenomenon, speciation. This is surprising in view of the special importance Darwin attached to extinction, and because the number of species extinctions in the history of life is almost the same as the number of originations; present-day biodiversity is the result of a trivial surplus of originations, cumulated over millions of years. For an evolutionary biologist to ignore extinction is probably as foolhardy as for a demographer to ignore mortality. The past decade has seen a resurgence of interest in extinction, yet research on the topic is still at a reconnaissance level, and our present understanding of its role in evolution is weak. Despite uncertainties, extinction probably contains three important elements. (i) For geographically widespread species, extinction is likely only if the killing stress is one so rare as to be beyond the experience of the species, and thus outside the reach of natural selection. (ii) The largest mass extinctions produce major restructuring of the biosphere wherein some successful groups are eliminated, allowing previously minor groups to expand and diversify. (iii) Except for a few cases, there is little evidence that extinction is selective in the positive sense argued by Darwin. It has generally been impossible to predict, before the fact, which species will be victims of an extinction event.


PNAS 1994. 91:6758-6763.


Background vs. Global Mass Extinction Events


- Background Rates = 96% of all extinctions.
- Phanerozoic average @ ~ 25% per 1 Myr.
- Geographic range & larval life styles regarding survivorship.
- Selective opportunities for diversification.
- The Red Queen Hypothesis.


Distribution of extinction intensities across Phanerozoic yields a mean of 25% per 1 Myr, which is the reciprocal of the mean species duration of ~4 Myr.




























The history of diversity of the three “evolutionary faunas” in the marine fossil record




Diversity-Dependent Competition
Is there a target equilibrium?


Summation of diversity of the three “evolutionary faunas” in the marine fossil record



Taxonomic survivorship curves

(A) Hypothetical curves






~Constant Probability of Extinction

- Red Queen Hypothesis: Must keep running to stay in same place.
- Constant evolution by competitors, predators, and parasites.

The BIG FIVE of the Phanerozoic

<u>Extinction Episode</u>	<u>Age, Ma</u>	<u>% Extinction</u>
• Cretaceous (K/T Boundary)	65 Ma	76
• Triassic	215 Ma	76
• Permian (P/T Boundary)	250 Ma	96
• Devonian	365 Ma	82
• Ordovician	440 Ma	85



The Permo-Triassic Boundary: The Mother of all Mass Extinctions

- Came close to losing all multicellular life.
- Considered one of the four major advancements.
- Box score of exterminations: 96% of all spp. & 50% of all families.
- Selectivity of the P/T Boundary.
- Multiple Causation Hypothesis.

The Four Major Advancements in Evolutionary Biology

- Origin of life
- Origin of multicellular life (Eukarya)
- Cambrian Explosion
- P/T Boundary Mass Extinction Event

P/T box score of extirminations

Rem: 96% of all spp. & 50% of all families


- 8 of 27 insects
- 21 of 27 reptiles
- 6 of 9 amphibians
- 70% of marine invertebrate genera including most corals.
- 1 major order of forams (the only time this has happened!).

Selectivity of the P/T Boundary


- 35% cosmopolitan genera vs. 93% endemic genera went extinct (same pattern as background extinction).
- END of the Trilobites as opposed to other marine arthropods.

Multiple Causation Hypothesis aka “World-went-to-hell” hypothesis

- Researchers tend to search for a single unified cause.....climate change, sea level change, oceanic anoxia, flood basalts, acid rain, etc.
- As much as 5 Ma separation period!
- Bolide impact hypothesis?








Impact Event at the Permian-Triassic Boundary: Evidence from Extraterrestrial Noble Gases in Fullerenes

Luann Becker,^{1*} Robert J. Poreda,² Andrew G. Hunt,² Theodore E. Bunch,³ Michael Rampino⁴


The Permian-Triassic boundary (PTB) event, which occurred about 251.4 million years ago, is marked by the most severe mass extinction in the geologic record. Recent studies of some PTB sites indicate that the extinctions occurred very abruptly, consistent with a catastrophic, possibly extraterrestrial, cause. Fullerenes (C₆₀ to C₂₄₀) from sediments at the PTB contain trapped helium and argon with isotope ratios similar to the planetary component of carbonaceous chondrites. These data imply that an impact event (asteroidal or cometary) accompanied the extinction, as was the case for the Cretaceous-Tertiary extinction event about 65 million years ago.

Nature 2001. 291:1530-1533.


Probable **competitive displacement** of brachiopods by bivalves




Probable **competitive displacement** of brachiopods by bivalves



Models of (A) **competitive displacement** and (B) **incumbent replacement**





Echinoid diversity increased during the Mesozoic and Cenozoic





K/T Boundary – Impact extinction

- 60 to 80% kill of all spp.
- END of the **Ammonites** as opposed to other marine molluscs.
- Bivalves were less selectively hit, broad range survival.
- Sea Urchins were selectively hit as well.










Bird extinctions on oceanic islands

60 species extinct in last 1500 yrs in Hawaii, following human colonization.

An estimated 2000 species of flightless rail used to live on islands in the Pacific...
...only 4 species remain today.

Approximately 1/5 of all bird species in the world have gone extinct in association with human colonization of the Pacific islands.



Current estimated extinction rates vs. the Big Five

- Based on rates of deforestation and patterns of tropical diversity and endemism ...in the next 30 years, 5-10% of Earth's species will go extinct.
- Such a rate is 100 to 1000 times the background extinction rate.
- How about when global warming kicks into high gear?
- Given the current rate of human population growth, this rate is likely to continue long enough to result in a mass extinction.
- **This will be the first mass extinction caused by an organism.**

The effect of magnetic fields on γ -ray bursts inferred from multi-wavelength observations of the burst of 23 January 1999

T. J. Galama¹, M. S. Briggs², R. A. M. J. Wijers³, P. M. Vreeswijk¹, E. Ro¹, D. Band⁴, J. van Paradijs^{1,5}, C. Kouveliotou⁶, R. D. Preece⁷, M. Bremer⁸, I. A. Smith⁹, R. P. J. Tilanus¹⁰, A. G. de Bruyn¹¹, R. G. Strom¹², G. Pooley¹³, A. J. Castro-Tirado¹⁴, M. Tavakoli¹⁵, C. Robinson¹⁶, K. Hurley¹⁷, J. Heise¹⁸, J. Telting¹⁹, R. G. M. Rutten²⁰, C. Packham²¹, R. Swaters¹, J. K. Davies²², A. Fassia²³, S. F. Green²⁴, M. J. Foster²⁵, R. Sagar²⁶, A. K. Pandey²⁷, Nilakshi²⁸, R. K. S. Yadav²⁹, E. O. Ofek³⁰, E. Leibowitz³¹, P. Ibbetson³², J. Rhoads³³, E. Falco³⁴, C. Peiry³⁵, C. Impey³⁶, T. R. Geballe³⁷ & D. Bhattacharya³⁸

Gamma-ray bursts (GRBs) are thought to arise when an extremely relativistic outflow of particles from a massive explosion (the nature of which is still unclear) interacts with material surrounding the site of the explosion. Observations of the evolving changes in emission at many wavelengths allow us to investigate the origin of the photons, and so potentially determine the nature of the explosion. Here we report the results of γ -ray, optical, infrared, submillimetre, millimetre and radio observations of the burst GRB990123 and its afterglow. Our interpretation of the data indicates that the initial and afterglow emissions are associated with three distinct regions in the fireball. The peak flux of the afterglow, one day after the burst, has a lower frequency than observed for other bursts; this explains the short-lived radio emission. We suggest that the differences between bursts reflect variations in the magnetic-field strength in the afterglow-emitting regions.

Nature 1999, 398:394-399.

Cosmic Explosion - Gamma-ray bursts are the most powerful explosions known in the Universe